Dark Energy and the Preposterous Universe

Sean Carroll, University of Chicago

Observation tells us: there is a dark, smooth, slowly–varying energy component comprising 70% of the energy in the universe.

- $\Omega_{clustered} = 0.3$; a flat universe

  implies ρ_{Dark} is smooth

- $\dot{a}^2 = \frac{8\pi G}{3} a^2 \rho - k$; acceleration

  implies ρ_{Dark} is slowly–varying



[Jaffe et al.]

Simplest candidate: vacuum energy, $\rho_{vac}(t, \vec{x}) = const.$

In fact, we have: $\rho_{vac} \approx 10^{-8} ergs/cm^3 \approx (10^{-3} eV)^4$.

This raises (at least) two big problems:

- Why is $\rho_{vac} \ll (10^{18} GeV)^4$?
- Why is $\rho_{vac} \sim \rho_{Matter}$?

Problem One: Why is the vacuum energy so small?

Vacuum energy is the most natural thing in the world.

Consider a single scalar field: $\rho_\phi = \frac{1}{2}\dot{\phi}^2 + \frac{1}{2}(\nabla\phi)^2 + V(\phi)$

$$\phi = const \longrightarrow \rho_\phi = V(\phi) = const.$$

Crucial point:
no known (unbroken) symmetry would prefer $V(\phi) = 0$.



How big a problem is the cosmological constant problem?

$$\rho_{vac} = \rho_{Higgs} + \rho_{QCD} + \rho_{0-point} + \rho_{\Lambda_0} + \ldots$$

$$(10^2 GeV)^4 \qquad (10^{-1} GeV)^4 \qquad \infty \qquad (10^{18} GeV)^4$$

So we predict: $\boxed{\rho_{vac}^{(theory)} \approx (10^{18} GeV)^4 = 10^{120} \rho_{vac}^{(obs)}}$

That's a cheat; really $\rho_{vac} = M_{vac}^4$; $\boxed{M_{vac}^{(theory)} = 10^{30} M_{vac}^{(obs)}}$

In fact, in supersymmetry, $M_{vac} \approx M_{\cancel{SUSY}} \approx 10^3 GeV = 10^{12} eV$,

so $M_{vac}^{(SUSY)} = 10^{15} M_{vac}^{(obs)}$.